

LOCALIZATION AND PATTERN FORMATION IN WIGNER REPRESENTATION VIA MULTIRESOLUTION

Antonina N. Fedorova, Michael G. Zeitlin

IPME RAS, St. Petersburg, V.O. Bolshoj pr., 61, 199178, Russia
e-mail: zeitlin@math.ipme.ru
e-mail: anton@math.ipme.ru
<http://www.ipme.ru/zeitlin.html>
<http://www.ipme.nw.ru/zeitlin.html>

We present an application of variational-wavelet analysis to quasiclassical calculations of solutions of Wigner equations related to nonlinear (polynomial) dynamical problems. (Naive) deformation quantization, multiresolution representations and variational approach are the key points. Numerical calculations demonstrates pattern formation from localized eigenmodes and transition from chaotic to localized (wavelet) types of behaviour.

Presented at VIII International Workshop on
Advanced Computing and Analysis Techniques in Physics Research,
Section III "Simulations and Computations in
Theoretical Physics and Phenomenology"
ACAT'2002, June 24-28, 2002, Moscow

Localization and pattern formation in Wigner representation via multiresolution

A. N. Fedorova and M. G. Zeitlin ^a

^aIPME, RAS, V.O. Bolshoj pr., 61, 199178, St. Petersburg, Russia

E-mail: zeitlin@math.ipme.ru, <http://www.ipme.ru/zeitlin.html>, <http://www.ipme.nw.ru/zeitlin.html>

We present an application of variational-wavelet analysis to quasiclassical calculations of solutions of Wigner equations related to nonlinear (polynomial) dynamical problems. (Naive) deformation quantization, multiresolution representations and variational approach are the key points. Numerical calculations demonstrates pattern formation from localized eigenmodes and transition from chaotic to localized (wavelet) types of behaviour.

In this paper we consider the applications of numerical-analytical technique based on local nonlinear harmonic analysis (wavelet analysis) to quasiclassical quantization of nonlinear (polynomial/rational) dynamical problems which appear in many areas of physics. Our starting point is the general point of view of deformation quantization approach at least on naive Moyal level [1]. So, for quantum calculations we need to find an associative (but non-commutative) star product $*$ on the space of formal power series in $\hbar$ with coefficients in the space of smooth functions such that $f * g = fg + \hbar\{f, g\} + \sum_{n \geq 2} \hbar^n B_n(f, g)$. In this paper we consider calculations of Wigner functions $W(p, q, t)$ (WF) as the solution of Wigner equation [1]

$$i\hbar \frac{\partial}{\partial t} W = H * W - W * H \quad (1)$$

corresponding to polynomial Hamiltonians $H(p, q, t)$. In general, equation (1) is nonlocal (pseudodifferential) for arbitrary Hamiltonians but in case of polynomial Hamiltonians we have only a finite number of terms in the corresponding series. For example, in stationary case after Weyl-Wigner mapping we have the following equation on WF in c-numbers [1]:

$$\left(\frac{p^2}{2m} + \frac{\hbar}{2i} \frac{p}{m} \frac{\partial}{\partial q} - \frac{\hbar^2}{8m} \frac{\partial^2}{\partial q^2} \right) W(p, q) + \quad (2)$$

$$U\left(q - \frac{\hbar}{2i} \frac{\partial}{\partial p}\right) W(p, q) = \epsilon W(p, q)$$

and after expanding the potential U into the Tay-

lor series we have two real finite partial differential equations. Our approach is based on extension of our variational-wavelet approach [2], [3]. Wavelet analysis is some set of mathematical methods, which gives us the possibility to work with well-localized bases in functional spaces and gives maximum sparse forms for the general type of operators (differential, integral, pseudodifferential) in such bases. We decompose the solutions of (1), (2) according to underlying hidden scales [2], [3] as

$$W(t, q, p) = \bigoplus_{i=i_c}^{\infty} \delta^i W(t, q, p) \quad (3)$$

where value i_c corresponds to the coarsest level of resolution c in the full multiresolution decomposition $V_c \subset V_{c+1} \subset V_{c+2} \subset \dots$. As a result the solution of any Wigner-like equations has the following multiscale/multiresolution decomposition via nonlinear high-localized eigenmodes, which corresponds to the full multiresolution expansion in all underlying scales starting from coarsest one (polynomial tensor bases are introduced in [3]; $\mathbf{x} = (x, y, p_x, p_y)$ e.g.):

$$W(t, \mathbf{x}) = \sum_{(i,j) \in \mathbb{Z}^2} a_{ij} \mathbf{U}^i \otimes V^j(t, \mathbf{x}), \quad (4)$$

$$V^j(t) = V_N^{j,slow}(t) + \sum_{l \geq N} V_l^j(\omega_l t), \quad \omega_l \sim 2^l$$

$$\mathbf{U}^i(\mathbf{x}) = \mathbf{U}_M^{i,slow}(\mathbf{x}) + \sum_{m \geq M} \mathbf{U}_m^i(k_m \mathbf{x}), \quad k_m \sim 2^m$$

Formula (4) gives the expansion into the slow




Figure 1. Localized mode contribution to Wigner function.




Figure 2. Chaotic-like Wigner function.

part and fast oscillating parts for arbitrary N , M . So, we may move from coarse scales of resolution to the finest one for obtaining more detailed information about our dynamical process. In contrast with different approaches formulae (3), (4) do not use perturbation technique or linearization procedures. So, by using wavelet bases with their best (phase) space/time localization properties we can describe localized (coherent) structures in quantum systems with complicated behaviour. Modeling demonstrates the appearance of stable patterns formation from high-localized coherent

structures or chaotic behaviour. Our (nonlinear) eigenmodes are more realistic for the modelling of nonlinear classical/quantum dynamical process than the corresponding linear gaussian-like coherent states. Here we mention only the best convergence properties of expansions based on wavelet packets, which realize the minimal Shannon entropy property and exponential control of convergence based on the norm introduced in [3]. Fig. 1 shows the high-localized eigenmode contribution to WF, while Fig. 2, 3 gives the representations for full solutions, constructed from the first 6 eigenmodes and demonstrate stable localized pattern formation (wavelet) and chaotic-like behaviour outside of KAM-like region. We can control the type of behaviour on the level of reduced algebraical variational system [3].




Figure 3. Localized pattern-like (wavelet) Wigner function.

REFERENCES

1. W. P. Schleich, Quantum Optics in Phase Space, Wiley, 2000
2. A.N. Fedorova, M.G. Zeitlin, Quantum Aspects of Beam Physics, 527-538, 539-550, World Scientific, 2002.
3. A.N. Fedorova, M.G. Zeitlin, arXiv preprints: physics/0206049, 0206050, 0206051, 0206052, 0206053, 0206054, 0212066, nlin/0206024.